

## Effectsof *Camellia sinensis* Ethanolic Extract on Histometric and Histopathological Healing Process of Burn Wound in Rat

<sup>1</sup>Mehrdad Karimi, <sup>2</sup>Pouya Parsaei, <sup>3</sup>Sayyed Yazdan Asadi, <sup>3</sup>Sareh Ezzati,  
<sup>2</sup>Reza Khadivi Boroujeni, <sup>4</sup>Alaleh Zamiri and <sup>5</sup>Mahmoud Rafieian-Kopaei

<sup>1</sup>Department of Surgery, Shahrekord University of Medical Sciences, Shahrekord, Iran

<sup>2</sup>Member of Young Researchers Club, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran

<sup>3</sup>Medical Plants Research Center, Faculty of Medicine,

Shahrekord University of Medical Sciences, Shahrekord, Iran

<sup>4</sup>Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran

<sup>5</sup>Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

**Abstract:** Green tea (*Camellia sinensis*) that were used as an ancient herbal drug in Asia, with having anti-oxidant and anti-inflammatory properties, may can improve wound healing process, therefore the present study were design to determinate the effect of green tea ethanolic extract on burn wound healing process in rat. For this experimental study, 36 healthy male Wistar rats were randomly selected into three groups of A, B and C which, respectively treated with, Vaseline + 0.6% green tea extract, Vaseline and normal saline for a 21 days period. Burn Wounds' areas were measured and calculated by caliper every other day and pathologic samples were taken at 3<sup>rd</sup>, 12<sup>th</sup> and 21<sup>st</sup> day for histopathological examinations. In the end, all data were analyzed by SPSS 16 using survival analysis (Breslow test), repeated measured ANOVA, one-way ANOVA and Mann-Whitney.  $P < 0.05$  was considered as statistically significant. Regarding Breslow test, the mean healing duration in group A with 18 days was shorter in compared to group B with 20.33 days, ( $P = 0.023$ ). In according to Mann-Whitney test, comparing groups A and C during the whole study, indicated no statistically significant differences in histopathological scores of epithelial regeneration and angiogenesis except inflammatory cells. Green tea extract could help wound healing process, probably effective on burn wounds healing.

**Key words:** Green Tea % *Camellia Sinensis* % Healing Process % Burn Wound

### INTRODUCTION

Herbal and natural products of folk medicine have been used for centuries in every culture all over the world. Data from WHO show that 70-80% of the world's population use herbal medicines as alternative medicine specially in middle east countries that is rich in herbal plant drugs like, Pakistan and Iran [1, 2].

Recently, the traditional use of plants for wound healing has received attention by the scientific community [3, 4]. Approximately one-third of all traditional medicines in use are for the treatment of wounds and skin disorders, compared to only 1-3% of modern drugs [5, 6].

Skin is one of the body's most important organs, which could prevent dehydration and influx of external factors [7].

Among all kinds of skin injuries, Burn wounds are very common skin injury in all communities, however, in developing countries burns consider as a major health problem because the incidence of severe complications is high and financial resources are limited. Normal wound healing response begins the moment the tissue is injured that comprises a complex pathophysiological process including several cellular and biochemical processes, e.g. inflammation, vessel formation and collagen deposition [8, 9]. Inflammation maintenance and inadequate vessel formation comprise the most noticeable causes of delayed wound healing [10, 11]. This delay may cause major wound infection that is responsible for 50-75% of hospital deaths. On the other hand wound fibrosis or abnormal accumulation of collagen around the wound could lead to an unpleasant scar that is very important in some parts of body like face and limbs [12, 8, 13].

Green tea (*Camellia sinensis*) were used as Asian people for health promotion healing diseases since 3000 B.C. This plant contains polyphenolic, in particular, catechin compounds [7, 13-17] Epicatechin, epicatechingallate, epigallocatechin and epigallocatechingallate that are responsible for anti-oxidant, anti-cancer, anti-aging and anti-inflammatory effects for this herbal medicinal plant [18]. Also it could prevent collagen production and accumulation and induce changes in immune responses in the healing phases [4, 5, 6, 18].

These compounds (e.g. epigallocatechingallate) have also been used as an agent for keratinocytes reproduction and distinction [14]. Also, its anti-fibrinogen effects have been confirmed in some animal models [15].

Due to previous studies, traditional experiences and easy accessibility, we investigated the effect of ethanolic extract of green tea on Burn wound healing process in wistar rat and determination of antioxidant Activity.

## MATERIALS AND METHODS

After obtaining approval in Shahrekord University of Medical Sciences (SKUMS) Ethics Committee by the number of 16.5.90, this experimental study was conducted in Medicinal Plants Research Center of SKUMS and Researches Centre of Medicinal Plants and Ethno-veterinary, Islamic azad university of shahrekord branch.

**Preparation of Extract:** Maceration method was employed to prepare the extract. For this purpose, 100g green tea (Herbarium No. 304, Medical Plants Research Center, SKUMS) was transported into an Erlenmeyer, 1 liter ethanol 70% (Nasr Co. Iran) was added and the solution was left at laboratory temperature. Forty eight hours later, the extract was filtrated through a filter paper and the pulp was squeezed to discharge. Then, the extract was concentrated by a rotary evaporator [17], dried and mixed with pure Vaseline (Ehsan Chemi, Iran) to make a Vaseline-based 0.6% ointment [19].

**Animals:** 36 Healthy male young Wistar Albino rats, weighing about (200-250g) obtained from animal house of Shahrekord university of medical sciences. They were

maintained under standard laboratory conditions of food and water and kept for quarantine for 24 hours. They were randomly assigned to three groups of A, B and C, the animals were treated with Vaseline+0.6% green tea extract (group A), Vaseline (group B) and normal saline (group C) for 21 days.

**Burn Wound Model:** After anaesthetizing Rats, by a combination of 20 mg/kg ketamine 10% (Alfasan Co., Netherlands) and 2mg/kg xylazine 2% (Alfasan Co., Netherlands), administered intramuscularly. Then, type II burning was created by putting 120°C Circular metal for 5 second on the shaved and scraped back of the rats [20]. Then rats were maintained for recovery in same situations and 12/12 hours light/dark cycle at 21±2°C.

**Treatment:** Performed every day at the same time by one person, treatment continued for 21 days, beginning from the day of making wound. For this, 1g Vaseline ointment containing green tea extract was topically applied on area of the burn wound in group A; the same was performed on group B and C using Vaseline ointment and normal saline (Daru-Pakhsh, Iran), respectively [21].

**Histometric Examinations:** Since the 2nd day of the study, the length of the remaining Burn wound areas was measured by a using a caliper and photographed every other day; each case's data was separately recorded [22, 21].

**Histopathological Examinations:** For pathologic sampling, 3 rats were randomly selected from groups at 3<sup>rd</sup>, 12<sup>th</sup> and 21<sup>st</sup> days of study period. After sacrifice them through anesthesia over dosage, the wound tissue histological specimens in full thickness accompanied with neighboring healthy skin were taken and fixed into 10% Neutral Buffered Formalin. Tissue processing was done by paraffin and transverse section, 5 micrometers thick, were made by means of Microtome fixed blade. The sections were stained by hematoxylin and eosin [23, 24]. Histopathological examinations, in view of inflammatory cells, epithelial regeneration and angiogenesis, were separately done by a pathologist who was blinded in grouping by scores that mentioned on Table 1.

Table 1: System for the histological features scores

Score	Epithelial Regeneration	Inflammatory cell	Angiogenesis
0	Absent	<5	Altered angiogenesis (1-2 vessels per site)
1	Starting	5 to 10	Few newly formed capillary vessels (3-6 vessels per site)
2	Incomplete (partial thickness)	11 to 20	Newly formed capillary vessels (7-10 vessels per site)
3	Complete (full thickness)	>20	Newly formed and well-structured capillary vessels (>10 vessels per site)

**Statistical Analysis:** Data analysis were done by SPSS 16 using survival analysis (Breslow test), repeated measures ANOVA, one-way ANOVA and Mann-Whitney.  $p < 0.05$  was significantly considered.

## RESULTS

**Morphological and Histometrical Findings:** Totally, 5 of 6 (83%) rats in group A (Vaseline+0.6% green tea extract) and 2 of 6 (33%) rats in group B (Vaseline) recovered by 21st day and no inflammation, eczema or infection were observed.

Regarding Breslow test, the mean healing duration in group A [18 days (SE=0.624)] was shorter compared to group B [20.33 days (SE = 0.385)] and it was statistically significant ( $P=0.023$ ).

The Burn wound area decrease process in group A, based on repeated measures ANOVA, was significantly faster than group B ( $p=0.002$ ). Diagram 1 indicates Burn

wound areas during the study and Table 2 indicates the mean decrease in burn area during 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> weeks.

Based on one-way ANOVA, the mean of increased burn area at the first week in group A were less than group B ( $P < 0.001$ ) on the other hand there was no significant difference in the mean decrease in burn area between groups A and B during 2<sup>nd</sup> week ( $P=0.08$ ) but during the 3<sup>rd</sup> week, the difference between the two groups was significant ( $P=0.013$ ).

It is notable that the decrease in burn area in group C was significantly less than the two other groups during the 21 days and there was no significant difference in the mean decrease in burn area between groups A and B during in all of the period ( $P=0.875$ ).

**Histopathological Findings:** Table 3 shows Mean Scores  $\pm$  Standard deviation of histopathological features of burn wound healing in different groups throughout the whole study.

Table 2: Comparison of mean decrease in burn area during the study (cm<sup>2</sup>)

Groups	1 <sup>st</sup> week	2 <sup>nd</sup> week	3 <sup>rd</sup> week	Total	Results of data analysis in 3 weeks
(Repeated measures) Vaseline+0.6% green tea extract(A)	-0.98	6.67	0.38	6.07	$p < 0.001$
Vaseline (B)	-3.28	7.75	1.40	5.87	$p < 0.001$
Normal saline(C)	-3.12	4.48	2.70	4.06	0.005
Results of data analysis (one-way ANOVA)	$< 0.001$	0.005	$< 0.001$	0.001	

Table 3: Histopathological evaluation of wound tissue samples at whole study

Wound healing' Scaling	Groups	Epithelial Regeneration	Inflammatory cells	Angiogenesis
Green tea+Vaseline (A)	0.72 $\pm$ 0.88		0.81 $\pm$ 1	3.317* $\pm$ 4.88
Vaseline (B)	0.67 $\pm$ 0.55	1.13 $\pm$ 1.66	1.69 $\pm$ 3	
Normal Saline (C)	0.72 $\pm$ 0.44	0.61 $\pm$ 2.22	1.02 $\pm$ 2.22	

-Values are mean  $\pm$  SD for the histological features in each group

-Asterisk in superscript indicates significant difference

-All values are significant at  $P < 0.05$




Diagram 1: Burn wound area (cm<sup>2</sup>) in all groups during the study




Fig. 1: Histopathologic sample Group A, appropriate angiogenesis and arranged collagen bands with low inflammatory cells. (H and E, X100, 400)




Fig. 2: Remaining some inflammatory cells (H and E, X100, 400)




Fig. 2: Incomplete epithelial in group C regeneration in group C (H and E, X100, 400)

In according to Mann-Whitney test, comparing groups A and C during the whole study, indicated no statistically significant differences in histopathological scores of epithelial regeneration ( $P=0.192$ ) and angiogenesis ( $P=0.106$ ) except inflammatory cells that was statistically significant ( $P=0.008$ ).

In addition, this comparison at 21<sup>st</sup> day of the study indicated statistically significant differences in no histopathological scores of epithelial regeneration, inflammatory cells except angiogenesis ( $P=0.361$ ,  $P=0.068$  and  $P=0.046$ , respectively).

Following microscopic photographs shows some difference between groups during the study.

## DISCUSSION

This study carried out to determine beneficial effects of green tea extract on healing of type II burn injuries in a rat model.

Wound healing is a very complex, multifactor sequence of events involving several cellular and biochemical processes [25]. It depends on the type and extent of damage, the general state of health and the tissue recovery ability.

Wound contracture is a process that occurs throughout the healing process, commencing in the fibroblastic stage whereby the area of the wound undergoes shrinkage [25, 26]. But some of problems that occurred may cause delay in wound healing that lead to increases the possibility of infection, inappropriate recovery and unpleasant scar [27, 26].

There are many studies in green tea healing aspects in diseases, like: anti-oxidant, anti-cancer, anti-aging, anti-inflammatory effects and preventive effects of collagen production and accumulation [28, 19]; the majority of these properties could be attributed to the plant's polyphenolic compound, i.e., catechin in the leaves [29, 30].

Consistent with our findings, Safari and Sadrzade's study indicated anti-oxidant effects of epigallocatechin, one of the green tea's compounds [31].

In addition, the beneficial effect of epicatechingallate and other phenolic components on wound healing quality and hence leaving more pleasant scar has been shown in Asadietal researches [20], that which confirm its effect on increased level of vascular endothelial growth factor and increasing angiogenesis in Group A in comparison to C in 3<sup>rd</sup> week by accelerated vessel formation and enhanced nitric oxide and cyclooxygenase [32]. Also Significant reduce, in number of inflammatory cells specially neutrophils in all duration of study in compare to control group C is the other results of the anti-oxidant components and phenolic substances and their activities [19, 30].

In morphometric examination, although the burn wound area were increased in all groups due to increasing the inflammation and skin tension on the dorsal region of rats at 1<sup>st</sup> week but it was significantly less than group B in A group ( $P<0.001$ ). In the other hand Analyzing the burn wounds areas during the study

demonstrated the significant effect of Vaseline+0.6% green tea extract in compare to control in decreasing the burn area in all period of study that is confirm the previous studies.

In according the properties mentioned for components of the green tea, it could help wound healing process. therefore, Further studies need to be done to identify and separate the group of active constituents responsible for anti-inflammatory activity and wound healing activity, Thus, after clinical trials in human, it can be considered as a natural, low cost easily available, treat to cure type II burns.

### ACKNOWLEDGEMENTS

This work was supported by the grant No. 626 by Research Department of SKUMS, Shahrekord, Iran. We also thank Medical Plants Research Center stuff, SKUMS, Researches Centre of Medicinal Plants and Ethno-veterinary, Islamic azad university of shahrekord branch, and all people who helped us throughout the study.

### REFERENCES

1. Divya, S., K. Naveen Krishna, S. Ramachandran and M.D. Dhanaraju, 2011. Wound Healing and *In vitro* Antioxidant Activities of Croton bonplandianum Leaf Extract in Rats. *Global Journal of Pharmacology*, 5(3): 159-163.
2. Marwat, S.K., M. Ajab Khan, M. Aslam Khan, M. Ahmad, M. Zafar, F. Rehman and S. Sultana, 2009. Fruit plant species mentioned in the Holy Qura'n and ahadith and their ethnomedicinal importance. *American-Eurasian Journal of Agricultural and Environmental Science*, 5(2): 284-295.
3. Kim, S.H., I.C. Kang and T.J. Yoon, 2001. Antitumor activities of a newly synthesized shikonin derivative, 2-hyim-DMNQS-33. *Cancer Letters*, 172: 171-175.
4. Ghorbani, A., 2005. Studies on pharmaceutical ethnobotany in the region of Turkman Sahra, north of Iran. *Journal of Ethnopharmacol.*, 102: 58-68.
5. Annan, K. and P.J. Houghton, 2007. Antibacterial, antioxidant and fibroblast growth stimulation of aqueous extracts of *Ficusasperifolia* Miq. and *Gossypiumarboreum* L., wound-healing plants of Ghana. *Journal of Ethnopharmacol.*, 119: 141-144.
6. Ghasemi Pirbalouti, A., S. Azizi and A. Koohpayeh, 2012. Healing potential of Iranian traditional medicinal plants on burn wounds in alloxan-induced diabetic rats. *Brazilian Journal of Pharmacognosy*, 22(2): 397-403.
7. McGrath, J.A. and S.M. Breathnach, 1998. Wound healing. In: R.H. Champion and J.L. Borton (eds.) *Rook's Textbook of Dermatology*. London Blackwell, pp: 336-55.
8. Mantle, D., M.A. Gok and T.W.J. Lennard, 2001. Adverse and beneficial effects of plant extracts on skin and skin disorders. *Adverse Drug Reaction and Toxicological Reviews*, 20: 89-103.
9. Mohajeri, D., M. Mesgari, Y. Doustar and M. Nazeri, 2011. Comparison of The Effect of Normal Saline and Silver Sulfadiazine on Healing of Skin Burn Wounds in Rats: A Histopathological Study. *Middle-East Journal of Scientific Research*, 10(1): 08-14.
10. Norrby, K., 1997. Angiogenesis: new aspects relating to its initiation and control. *APMIS*, 105(6): 417-37.
11. Mehdinezhad, B., A. Rezaei, D. Mohajeri and S. Safarmashaei, 2012. Comparison of *In vivo* Wound Healing Activity of Verbascum Thapsus flower extract with zinc oxide on experimental wound model in rabbits. *American. Eurasian Journal of Toxicological Sciences*, 4(1): 24-30.
12. Mirshamsi, M.H., J. Ayatollahi and M.H. Dashti-R, 2007. A Comparison Between Traumatic Wound Infections after Irrigating Them with Tap Water and Normal Saline. *World Journal of Medical Sciences*, 2(1): 58-61.
13. Hsu, S., 2005. Green tea and the skin. *Journal of American Academy of Dermatology*, 52(6): 1049-49.
14. Jankun, J., S.H. Selman, R. Swiercz and E. Skrzypczak-Jankun, 1997. Why drinking green tea could prevent cancer. *Nature*, 387(6633): 567.
15. Park, G., B.S. Yoon, J.H. Moon, B. Kim, E.K. Jun and S. Oh, 2008. Green tea polyphenol epigallocatechin-3-gallate suppresses collagen production and proliferation in keloid fibroblasts via inhibition of the STAT3-signaling pathway. *Journal of Investigative Dermatology. M.*, 128(10): 2429-41.
16. Hamaishi, K., R. Kojima and M. Ito, 2006. Anti-ulcer effect of tea catechin in rats. *Biol. Pharm. Bull.*, 29(11): 2206-13.
17. Khalaji, N. and T. Neyestani, 2007. The inhibitory effects of black and green teas (*Camellia sinensis*) on growth of pathogenic Escherichia coli, *in vitro*. *Iranian Journal of Nutrition Sciences*, 1(3): 33-38.

18. Kim, H.R., R. Rajaiah, S.R.W.Q. Satpute, M.T. Tan and J.E. Simon, 2008. Green tea protects rats against autoimmune arthritis by modulating disease-related immune events. *J. Nutrition*, 138(11): 2111-6.
19. Frei, B. and J.V. Higdon, 2003. Antioxidant activity of tea polyphenols *in vivo*: evidence from animal studies. *Journal of Nutrition*, 133(10): 3275-84.
20. Asadi, S.Y., A. Zamiri, S. Ezzati, P. Parsaei, M. Rafieian and H. Shirzad, 2011. Effect of alcoholic extract of green tea (*Camellia sinensis*) on the healing process in surgical and burn wounds in rats, *Journal of Birjand University of Medical Sciences*, 18(1): 1-9.
21. Baumann, L.S. and J. Spencer, 1999. The effects of topical vitamin E on the cosmetic appearance of scars. *Dermatol Surg*, 25: 34-5.
22. Cherly, S. Hedlund, 2007. Surgery of the Integumentary System. In: Theresa W. Fossum and Cheryl S. Hedlund. (eds.) *Small Animal Surgery*. 3<sup>rd</sup> ed. St.Louis: Mosby., pp: 159-76: 228-32.
23. Taghizadeh-Jahed, M., S.H. Jarolmasjed, S. Mohamadnejad and A. Rezaii, 2008. The effect of *Echinacea purpurea* aerial organ dried extract vs. Zinc oxide on skin wound healing in rat: amorphometric and histopathologic study. *J. Tehran University Medical*, 66(9): 625-632.
24. Derakhshanfar, A., M. Oloumi and M. Mirzaie, 2010. Study on the effect of *Peganum harmala* extract on experimental skin wound healing in rat: pathological and biomechanical findings. *Journal of Comparative Clinical Pathology*, 19(24): 169-172.
25. Philips, G.D., R.A. Whitehe and D.R. Kington, 1991. Initiation and pattern of angiogenesis in wound healing in the rat. *American Journal of Anatomy*, 192: 257-262.
26. Ghasemi Pirbalouti, A., S. Azizi and A. Koohpayeh, 2012. Healing potential of Iranian traditional medicinal plants on burn wounds in alloxan-induced diabetic rats. *Brazilian Journal of Pharmacognosy*, 22(2): 397-403.
27. Tredget, E.E., B. Nedelec, P.G. Scott and A. Ghahary, 1997. Hypertrophic scars, keloids and contractures: the cellular and molecular basis for therapy. *Surgical Clinics of North America*, 77(3): 701-30.
28. Nakamuta, M., N. Higashi, M. Kohjima, M. Fukushima, S. Ohta, K. Kotoh, N. Kobayashi and M. Enjoji, 2005. Epigallocatechin-3-gallate, a polyphenol component of green tea, suppresses both collagen production and collagenase activity in hepatic stellate cells. *International Journal of Molecular Medicine*, 16(4): 677-81.
29. Yen, G.C. and H.Y. Chen, 1998. Scavenging effect of various tea extracts on superoxide derived from the metabolism of mutagens. *Biosci. Biotech. Biochem.*, 62(9): 1768-70.
30. Katiyar, S.K. and C.A. Elmet, 2001. Green tea polyphenolic antioxidants and skin photoprotection (Review). *International Journal of Oncology*, 18(6): 1307-13.
31. Saffari, Y. and S.M.H., 2004. Sadrzadeh. Green tea metabolite EGCG protects membranes against oxidative damage *in vitro*. *Life Sciences*, 74(12): 1513-8.
32. Kapoor, M., R. Howard, I. Hall and I. Appleton, 2004. Effects of epicatechingallate on wound healing and scar formation in a full thickness incisional wound healing model in rats. *American Journal of Pathology*, 165(1): 299-307.